

Nadine Wehres¹, Frank Lewen¹, Christian Endres²,
Marius Hermanns¹, Stephan Schlemmer¹

HETERODYNE RECEIVER FOR LABORATORY SPECTROSCOPY OF MOLECULES OF ASTROPHYSICAL IMPORTANCE

¹Physikalisches Institut, University of Cologne


²Max Planck Institute for Extraterrestrial Physics, Garching



6/24/2016

71st ISMS Champaign-Urbana, IL




Emission Spectrometer






Heterodyne Receiver



- 75-110 GHz Frequency Range / ALMA 3 Band
- Room Temp. Low Noise Amplifier <390K @ 20dB Gain
- Mixer: LO 12.5 – 18.3 GHz @ x6
- IF Out: 6.35 GHz @ 2.5 GHz BW

IF Processor


- IN: 6.35 GHz @ 2.5 GHz BW
- Amplification & Filters
- Continuum Detector to set Amplification
- OUT: 0-2.5 GHz



X-FFTS





- IN: 0-2.5 GHz I/Q or Single
- 2.5 GHz BW @ 5 GS/s
- 10 bit ADC @ >55 dBc SFDR
- FFT with 32768 Channels @ 76.3 kHz spacing
- runs in APEX since 2009





More Information: MPIfR Bonn, Germany



Emission Spectrum of Methyl cyanide





Emission Spectrum of Ethanol




Emission Spectrum of Ethanol




Comparison Emission vs. Absorption




Just In





Methyl cyanide with copper cube absorption in 160 minutes





Chirped Pulse Setup



CP Spectrum of Methyl cyanide



CP Spectrum of Methyl cyanide




Team & Acknowledgements



Prof. Dr.
Schlemmer



Dr. Endres¹



Dr. Wehres



Dr. Lewen



Laboratory Astrophysics Group

CP: Oliver Zingsheim, Bettina Heyne

¹ present address: Max Planck Institute for Extraterrestrial Physics, Garching, Germany



